

98-99	1	1	2	8	3	56	4	405	5	100000
Individual	6	2401	7	9	8	36	9	11	10	9

98-99	1	3	2	-24	3	$\frac{1}{2}$	4	$\frac{1}{2}$	5	6
Group	6	12	7	4	8	7	9	12	10	135

# Individual Events

- 11 The circumference of a circle is  $14\pi$  cm. Let  $X$  cm be the length of an arc of the circle, which subtends an angle of  $\frac{1}{7}$  radian at the centre. Find the value of  $X$ .

Let  $r$  be the radius of the circle.  $2\pi r = 14\pi \Rightarrow r = 7$

$$X = r\theta = 7 \times \frac{1}{7} = 1$$


- 12 In Figure 1,  $ABCDEF$  is a regular hexagon with area equal to  $3\sqrt{3}$  cm<sup>2</sup>. Let  $X$  cm<sup>2</sup> be the area of the square  $PQRS$ , find the value of  $X$ .

Area of the hexagon =  $6 \times \text{areas of } \triangle AOB$

$$3\sqrt{3} = 6 \cdot \frac{1}{2} \cdot OB^2 \sin 60^\circ = \frac{3\sqrt{3}}{2} \cdot OB^2$$

$$OB^2 = 2$$

$$\text{Area of the square} = (2OB)^2 = 4 \times 2 = 8$$



- 13 8 points are given and no three of them are collinear. Find the number of triangles formed by using any 3 of the given points as vertices.

$$\text{The number of triangles formed} = {}_8C_3 = \frac{8 \times 7 \times 6}{1 \times 2 \times 3} = 56$$

- 14 In Figure 2, there is a  $3 \times 3$  square.


Let  $\angle a + \angle b + \dots + \angle i = X^\circ$ , find the value of  $X$ .

$$\angle c = \angle e = \angle g = 45^\circ$$

$$\angle a + \angle i = 90^\circ, \angle b + \angle f = 90^\circ, \angle d + \angle h = 90^\circ$$

$$\angle a + \angle b + \dots + \angle i = 45^\circ \times 3 + 90^\circ \times 3 = 405^\circ$$

$$X = 405$$



- 15 How many integers  $n$  are there between 0 and  $10^6$ , such that the unit digit of  $n^3$  is 1?

$1^3 = 1$ , the unit digit of  $n$  must be 1, there are  $10^6 \div 10 = 100000$  possible integers.

- 16 Given that  $a, b, c$  are positive integers and  $a < b < c = 100$ , find the number of triangles formed with sides equal  $a$  cm,  $b$  cm and  $c$  cm.

By triangle inequality:  $a + b > c = 100$

Possible pairs of  $(a, b)$ : (2, 99), (3, 98), (3, 99), (4, 97), (4, 98), (4, 99), ... ,

(50, 51), (50, 52), ... , (50, 99), .....

(98, 99)

$$\text{Total number of triangles} = 1 + 2 + \dots + 48 + 49 + 48 + \dots + 2 + 1$$

$$= \frac{1+49}{2} \times 49 \times 2 - 49 = 2401$$

- 17 A group of youngsters went for a picnic. They agreed to share all expenses. The total amount used was \$288. One youngster had no money to pay his share, and each of the others had to pay \$4 more to cover the expenses. How many youngsters were there in the group?

Let the number of youngsters be  $n$ .

$$\frac{288}{n-1} - \frac{288}{n} = 4$$

$$72 = n^2 - n, n = 9$$

- 18 A two-digit number is equal to 4 times the sum of the digits, and the number formed by reversing the digits exceeds 5 times the sum of the digits by 18. What is the number?

Let the unit digits of the original number be  $x$  and the tens digit by  $y$ .

$$10y + x = 4(x + y) \dots\dots\dots(1)$$

$$10x + y - 5(x + y) = 18 \dots\dots(2)$$

$$\text{From (1), } 6y = 3x \Rightarrow x = 2y \dots\dots\dots(3)$$

Sub. (3) into (2):  $20y + y - 5(2y + y) = 18 \Rightarrow y = 3, x = 6$ ; the number is 36.

- 19 Given that the denominator of the  $1001^{\text{th}}$  term of the following sequence is 46, find the

numerator of this term.  $\frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \dots$

Suppose the numerator of the  $1001^{\text{th}}$  term is  $n$ .

$$1 + 2 + 3 + \dots + 44 + n = 1001, n \leq 45$$

$$\frac{1}{2}(45)(44) + n = 1001, n = 1001 - 990 = 11$$

- 110 In the following addition, if the letter 'S' represents 4, what digit does the letter 'A' represent?

SEE

$$3E + 4 = 10a + Y \dots\dots\dots(1), \text{ where } a \text{ is the carry digit in the tens digit.}$$

4EE

SEE

$$4E + a = 10b + 4 \dots\dots\dots(2), \text{ where } b \text{ is the carry digit in the hundreds digit.}$$

4EE

SEE

$$4 \times 3 + Y + b = 10E + A \dots\dots\dots(3)$$

4EE

+ YES

From (3),  $E = 1$  or  $2$

+ YE4

EASY

When  $E = 1$ , (1)  $\Rightarrow Y = 7, a = 0$ , (2)  $\Rightarrow b = 0$ , (3)  $\Rightarrow A = 9$

When  $E = 2$ , (2)  $\Rightarrow a = 1, Y = 0$  reject because  $YE4$  is a 3-digit number.

EA4Y

$\therefore A = 9$

# Group Events

- G1 If  $a$  is a prime number and  $a^2 - 2a - 15 < 0$ , find the greatest value of  $a$ .

$(a + 3)(a - 5) < 0 \Rightarrow a < 5$ , the greatest prime number is 3.

- G2 If  $a : b : c = 3 : 4 : 5$  and  $a + b + c = 48$ , find the value of  $a - b - c$ .

$a = 3k, b = 4k, c = 5k$ ; sub. into  $a + b + c = 48 \Rightarrow 3k + 4k + 5k = 48 \Rightarrow k = 4$

$a = 12, b = 16, c = 20, a - b - c = 12 - 16 - 20 = -24$

- G3 Find the value of  $\log(\sqrt{3+\sqrt{5}} + \sqrt{3-\sqrt{5}})$ .

$$\begin{aligned} \log(\sqrt{3+\sqrt{5}} + \sqrt{3-\sqrt{5}}) &= \log\left(\sqrt{\frac{6+2\sqrt{5}}{2}} + \sqrt{\frac{6-2\sqrt{5}}{2}}\right) = \log\left(\frac{\sqrt{(1+\sqrt{5})^2} + \sqrt{(\sqrt{5}-1)^2}}{\sqrt{2}}\right) \\ &= \log\left(\frac{1+\sqrt{5}+\sqrt{5}-1}{\sqrt{2}}\right) = \log\left(\frac{2\sqrt{5}}{\sqrt{2}}\right) = \log(\sqrt{2}\sqrt{5}) = \log\sqrt{10} = \frac{1}{2} \end{aligned}$$

- G4 Find the area enclosed by the straight line  $x + 4y - 2 = 0$  and the two coordinate axes.

$x$ -intercept = 2,  $y$ -intercept =  $\frac{1}{2}$ ; the area =  $\frac{1}{2} \times \frac{1}{2} \times 2 = \frac{1}{2}$

- G5 Natural numbers are written in order starting from 1 until  $198^{\text{th}}$  digit as shown 123456789101112...... If the number obtained is divided by 9, find the remainder.

198 digits

123456789 has 9 digits, 10111213...9899 has  $90 \times 2 = 180$  digits

$\therefore$  1234567891011...9899100101102 has 198 digits.

$1+2+3+\dots+9 = 45$ ,  $11+12+\dots+19$  is also divisible by 9,...,  $91+92+\dots+99$  is divisible by 9.

$10+20+\dots+90$  is divisible by 9,  $\therefore$  the remainder is the same as 100101102 divided by 9.

$1+1+1+1+2 = 6$ , the remainder is 6.

- G6 The average of 2,  $a$ , 5,  $b$ , 8 is 6. If  $n$  is the average of  $a, 2a+1, 11, b, 2b+3$ , find the value of  $n$ .

$$2 + a + 5 + b + 8 = 30 \dots\dots\dots(1), a + 2a + 1 + 11 + b + 2b + 3 = 5n \dots\dots\dots(2)$$

From (1):  $a + b = 15$ ; (2)  $5n = 3a + 3b + 15 = 3(a + b) + 15 = 3 \times 15 + 15 = 60 \Rightarrow n = 12$

G7 If  $p = 2x^2 - 4xy + 5y^2 - 12y + 16$ , where  $x$  and  $y$  are real numbers, find the least value of  $p$ .

$$p = 2x^2 - 4xy + 2y^2 + 3y^2 - 12y + 16 = 2(x - y)^2 + 3(y^2 - 4y + 4) + 4 = 2(x - y)^2 + 3(y - 2)^2 + 4$$

$p \geq 4$ , the least value of  $p$  is 4.

G8 Find the unit digit of  $333^{335}$ .

$3^1 = 3, 3^2 = 9, 3^3 = 27, 3^4 = 81$ , the unit digit of  $3^{4m}$  is 1, where  $m$  is any positive integer.

$$333^{335} = 333^{4 \times 83 + 3} = (333^4)^{83} \times 333^3 = (\dots 1)^{83} \times (\dots 3^3) = \dots 7, \text{ the unit digit is } 7.$$

G9 In Figure 1,  $\angle MON = 20^\circ$ ,  $A$  is a point on  $OM$ ,  $OA = 4\sqrt{3}$ ,  $D$  is a point on  $ON$ ,  $OD = 8\sqrt{3}$ ,  $C$  is any point on  $AM$ ,  $B$  is any point on  $OD$ . If  $\ell = AB + BC + CD$ , find the least value of  $\ell$ .

Reflect the figure along the line  $OM$ , then reflect the  $O$  figure between  $\angle MON_1$  along the line  $ON_1$ .

$$\angle NOM_2 = 3 \times 20^\circ = 60^\circ$$


$$\ell = AB + BC + CD = AB_1 + B_1C + CD$$

$$\ell = A_2B_1 + B_1C + CD$$

$\ell$  is the shortest when  $A_2, B_1, C, D$  are collinear.

By cosine formula on  $\triangle OA_2D$ ,

$$\begin{aligned} \text{Shortest } \ell = A_2D &= \sqrt{(4\sqrt{3})^2 + (8\sqrt{3})^2 - 2(4\sqrt{3})(8\sqrt{3})\cos 60^\circ} \\ &= \sqrt{48 + 192 - 96} = 12 \end{aligned}$$



G10 In figure 2,  $P$  is a point inside the square  $ABCD$ ,  $PA = a$ ,  $PB = 2a$ ,  $PC = 3a$  ( $a > 0$ ). If  $\angle APB = x^\circ$ , find the value of  $x$ .  
Rotate  $\triangle APB$  by  $90^\circ$  in anti-clockwise direction about  $B$ .

Let  $P$  rotate to  $Q$ ,  $A$  rotate to  $E$ .

$\triangle APB \cong \triangle EQB$  (by construction)

$EQ = a$ ,  $BQ = 2a = PB$ . Join  $AQ$ .

$\angle PBQ = 90^\circ$  (Rotation)

$\angle ABQ = 90^\circ - \angle ABP = \angle PBC$

$AB = BC$  (sides of a square)

$\triangle ABQ \cong \triangle CBP$  (SAS)

$AQ = CP = 3a$  (corr. sides  $\cong \triangle$ 's)

$\therefore \angle PBQ = 90^\circ$  (Rotation)

$$\begin{aligned} \therefore PQ^2 &= PB^2 + QB^2 \text{ (Pyth. Theorem)} \\ &= (2a)^2 + (2a)^2 = 8a^2 \end{aligned}$$

$$AP^2 + PQ^2 = a^2 + 8a^2 = 9a^2$$

$$AQ^2 = (3a)^2$$

$$\therefore AP^2 + PQ^2 = AQ^2$$

$\angle APQ = 90^\circ$

$\therefore \angle PBQ = 90^\circ$  and  $PB = QB$

$\therefore \angle BPQ = 45^\circ$

$$\angle APB = 45^\circ + 90^\circ = 135^\circ$$

