

07-08 Individual	1	1.8	2	96	3	64	4	$\frac{12}{61}$	5	300
	6	27	7	3	8	$\frac{2007}{1004}$	9	2	10	-3

07-08 Group	1	1	2	34891	3	$\frac{8}{9}$	4	48	5	$\frac{1}{2}$
	6	1	7	$\frac{25}{12}$	8	3	9	6	10	$\frac{4\sqrt{41}}{5}$

## Individual Events

- 11 In Figure 1,  $ABC$  is a triangle,  $AB = 13$  cm,  $BC = 14$  cm and  $AC = 15$  cm.  $D$  is a point on  $AC$  such that  $BD \perp AC$ . If  $CD$  is longer than  $AD$  by  $X$  cm, find the value of  $X$ .

$$X = CD - AD = BC \cos C - AB \cos A$$

$$X = 14 \cdot \frac{15^2 + 14^2 - 13^2}{2 \cdot 15 \cdot 14} - 13 \cdot \frac{15^2 + 13^2 - 14^2}{2 \cdot 15 \cdot 13} = 2 \cdot \frac{14^2 - 13^2}{30}$$

$$X = \frac{27}{15} = \frac{9}{5}$$


Method 2

$$BD^2 = 14^2 - CD^2 = 13^2 - AD^2$$

$$14^2 - 13^2 = CD^2 - AD^2$$

$$(14 + 13)(14 - 13) = (CD + AD)(CD - AD)$$

$$27 = 15(CD - AD) \Rightarrow X = \frac{27}{15} = \frac{9}{5}$$



- 12 Given that a trapezium  $PQRS$  with dimensions  $PQ = 6$  cm,  $QR = 15$  cm,  $RS = 8$  cm and  $SP = 25$  cm, also  $QR \parallel PS$ . If the area of  $PQRS$  is  $Y$  cm<sup>2</sup>, find the value of  $Y$ .

Let the height of the trapezium be  $h$  cm ( $= QW$ ).From  $Q$ , draw  $QT \parallel RS$ , which intersect  $PS$  at  $T$ . $QRST$  is a // -gram (2 pairs of // -lines) $TS = 15$  cm,  $QT = 8$  cm (opp. sides, // -gram)


$$PT = (25 - 15) \text{ cm} = 10 \text{ cm}$$

$$QP^2 + QT^2 = PT^2$$

 $\angle PQT = 90^\circ$  (Converse, Pyth. Theorem)

$$\frac{1}{2} 6 \times 8 = \text{area of } \triangle PQT = \frac{1}{2} 10 \times h, h = 4.8$$

$$\text{Area of the trapezium} = Y = \frac{15 + 25}{2} \times 4.8 = 96$$



- 13 Given that  $x_0$  and  $y_0$  are positive integers satisfying the equation  $\frac{1}{x} + \frac{1}{y} = \frac{1}{15}$ . If  $35 < y_0 < 50$

and  $x_0 + y_0 = z_0$ , find the value of  $z_0$ .

$$15(x + y) = xy$$

$$xy - 15xy - 15y + 225 = 225$$

$$(x - 15)(y - 15) = 225$$

$$(x - 15, y - 15) = (1, 225), (3, 75), (5, 45), (9, 25), (15, 15), (25, 9), (45, 5), (75, 3), (225, 1)$$

$$\therefore 35 < y_0 < 50, y_0 - 15 = 25, x_0 - 15 = 9$$

$$y_0 = 40 \text{ and } x_0 = 24; z_0 = 24 + 40 = 64$$

- I4 Let  $a, b, c$  and  $d$  be real numbers. If  $\frac{a}{b} = \frac{1}{2}$ ,  $\frac{b}{c} = \frac{3}{2}$ ,  $\frac{c}{d} = \frac{4}{5}$  and  $\frac{ac}{b^2 + d^2} = e$ , find the value of  $e$ .

$$b = 2a; \quad \frac{a}{b} \times \frac{b}{c} = \frac{1}{2} \times \frac{3}{2} \Rightarrow c = \frac{4a}{3}; \quad \frac{a}{b} \times \frac{b}{c} \times \frac{c}{d} = \frac{1}{2} \times \frac{3}{2} \times \frac{4}{5} \Rightarrow d = \frac{5a}{3}$$

$$e = \frac{ac}{b^2 + d^2} = \frac{a \times \frac{4a}{3}}{(2a)^2 + \left(\frac{5a}{3}\right)^2} = \frac{\frac{4}{3}}{4 + \frac{25}{9}} = \frac{4}{3} \times \frac{9}{61} = \frac{12}{61}$$

Method 2  $a : b = 1 : 2, b : c = 3 : 2$

$a : b : c$

$1 : 2$

$\frac{3 : 2}{3 : 6 : 4}$

$3 : 6 : 4$

$\therefore a : b : c : d = 3 : 6 : 4 : 5$

Let  $a = 3k, b = 6k, c = 4k, d = 5k$

$$e = \frac{ac}{b^2 + d^2} = \frac{(3k)(4k)}{(6k)^2 + (5k)^2} = \frac{12}{61}$$

- I5 In Figure 2, the large rectangle is formed by eight identical small rectangles. Given that the length of the shorter side of the large rectangle is 40 cm and the area of the small rectangle is  $A \text{ cm}^2$ , find the value of  $A$ .



Let the length of a small rectangle be  $y \text{ cm}$ ;

the width of a small rectangle be  $x \text{ cm}$ .

Then  $x + y = 40, y = 3x$

$x = 10, y = 30$

$A = xy = 300$



- I6 In Figure 3,  $\triangle ABC$  is an equilateral triangle. It is formed by several identical equilateral triangles. If there are altogether  $N$  equilateral triangles in the figure, find the value of  $N$ .


Number of equilateral triangles of side 1 is 16.

Number of equilateral triangles of side 2 is 7.

Number of equilateral triangles of side 3 is 3.

Number of equilateral triangles of side 4 is 1.

Total number of equilateral triangles is 27.



- I7 Let  $r$  be the larger real root of the equation  $\frac{4}{y+1} + \frac{5}{y-5} = -\frac{3}{2}$ . Find the value of  $r$ .

$$8(y-5) + 10(y+1) = -3(y^2 - 4y - 5)$$

$$3y^2 + 6y - 45 = 0$$

$$y^2 + 2y - 15 = 0$$

$$(y+5)(y-3) = 0$$

$$y = -5 \text{ or } 3; r = 3$$

I8 Let  $x$  be a rational number and  $w = \left| x + \frac{2007}{2008} \right| + \left| x - \frac{2007}{2008} \right|$ . Find the smallest possible value of  $w$ .

$$\text{If } x < -\frac{2007}{2008}, w = -x - \frac{2007}{2008} - x + \frac{2007}{2008} = -2x, \therefore w = -2x > \frac{2007}{1004}$$

$$\text{If } -\frac{2007}{2008} \leq x < \frac{2007}{2008}, w = x + \frac{2007}{2008} - x + \frac{2007}{2008} = \frac{2007}{1004}, \therefore w = \frac{2007}{1004}$$

$$\text{If } \frac{2007}{2008} \leq x, w = x + \frac{2007}{2008} + x - \frac{2007}{2008} = 2x, \therefore w = 2x \geq \frac{2007}{1004}$$

The smallest possible value of  $w$  is  $\frac{2007}{1004}$ .

I9 Let  $m$  and  $n$  be positive integers. Given that the number 2 appears  $m$  times and the number 4 appears  $n$  times in the expansion  $\left( \left( \left( (2)^2 \right)^{\cdot} \right)^{\cdot} \right)^2 = \left( \left( \left( (4)^4 \right)^{\cdot} \right)^{\cdot} \right)^4$ . If  $k = \frac{m}{n}$ , find the value of  $k$ .

$$2^{(2^{m-1})} = 4^{(4^{n-1})}$$

$$2^{(2^{m-1})} = 2^{2(2^{2n-2})}$$

$$2^{m-1} = 2^{2n-1} \Rightarrow m = 2n$$

$$k = 2$$

I10 Find the value of  $\log_2(\sin^2 45^\circ) + \log_2(\cos^2 60^\circ) + \log_2(\tan^2 45^\circ)$ .

$$\log_2 \frac{1}{2} + \log_2 \frac{1}{4} + \log_2 1 = -1 - 2 + 0 = -3$$

Group Events

- G1 Given that the decimal part of  $5 + \sqrt{11}$  is  $A$  and the decimal part of  $5 - \sqrt{11}$  is  $B$ . Let  $C = A + B$ , find the value of  $C$ .

$$3 < \sqrt{11} < 4; 5 + \sqrt{11} = 5 + 3 + A = 8 + A; 5 - \sqrt{11} = 5 - (3 + A) = 2 - A = 1 + (1 - A) \\ B = 1 - A; A + B = 1; C = 1$$

- G2 A total number of  $x$  candies,  $x$  is a positive integer, can be evenly distributed to 851 people as well as 943 people. Find the least possible value of  $x$ .

$$851 = 23 \times 37; 943 = 23 \times 41$$

$$x = 23 \times 37 \times m = 23 \times 41 \times n, \text{ where } m, n \text{ are positive integers.}$$

$$37m = 41n$$

$$\therefore 37, 41 \text{ are relatively prime, the minimum } m = 41$$

$$\text{The least possible value of } x = 23 \times 37 \times 41 = 851 \times 41 = 34891$$

- G3 In Figure 1,  $ABCD$  is a tetrahedron with side length of 2 cm. If the

volume of the tetrahedron is  $\sqrt{R} \text{ cm}^3$ , find the value of  $R$ .

Let  $M$  be the mid point of  $BC$ . ( $BM = MC = 1 \text{ cm}$ )

$$\triangle ABM \cong \triangle ACM \text{ (SSS)}$$

$$AM = \sqrt{3} \text{ cm (Pythagoras Theorem)}$$


$$O \text{ is the centroid of } \triangle ABC. AO = \frac{2\sqrt{3}}{3} \text{ cm}$$

$$DO = \text{height of the tetrahedron} = \sqrt{AD^2 - AO^2} = \sqrt{\frac{8}{3}} \text{ cm}$$

$$\text{Volume} = \sqrt{R} \text{ cm}^3 = \frac{1}{3} \cdot \frac{1}{2} \cdot 2^2 \sin 60^\circ \sqrt{\frac{8}{3}} \text{ cm}^3$$

$$\sqrt{R} = \frac{1}{3} \sqrt{8}$$

$$R = \frac{8}{9}$$



- G4 Given that  $x$  is a positive integer and  $x < 60$ . If  $x$  has exactly 10 positive factors, find the value of  $x$ .

Note that except for the perfect square numbers (say 25), all positive integers have even numbers of factors. For a number  $x < 60$  which has 10 positive factors,  $x$  will be divisible by as many numbers  $< 8$  as possible. One possible choice would be 48. The positive factors are 1, 2, 3, 4, 6, 8, 12, 16, 24 and 48.

Method 2 Express  $N$  as unique prime factorization:  $p_1^{r_1} p_2^{r_2} \cdots p_n^{r_n}$ , then the number of factors is  $(1 + r_1)(1 + r_2) \cdots (1 + r_n) = 10 = 2 \times 5 \Rightarrow n = 2, 1 + r_1 = 2, 1 + r_2 = 5 \Rightarrow r_1 = 1, r_2 = 4$   
For prime numbers 2, 3, 5, ...,  $2^4 \times 3 = 48 < 60$ , other combinations will exceed 60.

- G5 Given that  $90^\circ < \theta < 180^\circ$  and  $\sin \theta = \frac{\sqrt{3}}{2}$ . If  $A = \cos(180^\circ - \theta)$ , find the value of  $A$ .

$$\theta = 120^\circ, A = \cos 60^\circ = \frac{1}{2}$$

- G6 Let  $x$  be a positive real number. Find the minimum value of  $x^{2008} - x^{1004} + \frac{1}{x^{1004}}$ .

Let  $t = x^{1004}$ , then  $t^2 = x^{2008}$

$$\begin{aligned} x^{2008} - x^{1004} + \frac{1}{x^{1004}} &= t^2 - t + \frac{1}{t} = t^2 - 1 - \frac{t^2 - 1}{t} + 1 \\ &= (t^2 - 1) \left( 1 - \frac{1}{t} \right) + 1 = \frac{(t^2 - 1)(t - 1)}{t} + 1 = \frac{(t - 1)^2(t + 1)}{t} + 1 \end{aligned}$$

Clearly  $t = x^{1004} > 0$ ,  $(t - 1)^2(t + 1) \geq 0$ ,  $\frac{(t - 1)^2(t + 1)}{t} + 1 \geq 0 + 1 = 1$ , equality holds when  $t = 1$ .

$\therefore$  When  $x = 1$ , the minimum value of  $x^{2008} - x^{1004} + \frac{1}{x^{1004}}$  is 1.

- G7 Let  $x$  and  $y$  be real numbers satisfying  $\begin{cases} \left(x - \frac{1}{3}\right)^3 + 2008\left(x - \frac{1}{3}\right) = -5 \\ \left(y - \frac{7}{4}\right)^3 + 2008\left(y - \frac{7}{4}\right) = 5 \end{cases}$ . If  $z = x + y$ , find the

value of  $z$ .

Let  $a = x - \frac{1}{3}$ ,  $b = y - \frac{7}{4}$ . Add up the two equations:  $a^3 + b^3 + 2008(a + b) = 0$

$$(a + b)(a^2 - ab + b^2) + 2008(a + b) = 0$$

$$(a + b)(a^2 - ab + b^2 + 2008) = 0$$

$$a + b = 0 \text{ or } a^2 - ab + b^2 + 2008 = 0$$

$$\text{But } a^2 - ab + b^2 + 2008 = \left(a - \frac{b}{2}\right)^2 + \frac{b^2}{4} + 2008 \geq 2008 \neq 0$$

$$\therefore a + b = x - \frac{1}{3} + y - \frac{7}{4} = 0; z = x + y = \frac{25}{12}$$

- G8 Let  $R$  be the remainder of  $1 \times 3 \times 5 \times 7 \times 9 \times 11 \times 13 \times 15 \times 17 \times 19$  divided by 4. Find the value of  $R$ .  
Note that if  $N$  and  $m$  are positive integers,  $0 \leq m \leq 99$  and  $x = 100N + m$ , then the remainder when  $x$  divided by 4 is the same as that when  $m$  is divided by 4.

$$\begin{aligned} \text{Product} &= 1 \times (4-1) \times (4+1) \times (4 \times 2 - 1) \times (4 \times 2 + 1) \times (4 \times 3 - 1) \times (4 \times 3 + 1) \times (4 \times 4 - 1) \times (4 \times 4 + 1) \times (4 \times 5 - 1) \\ &= 1 \times (4+1) \times (4 \times 2 + 1) \times (4 \times 3 + 1) \times (4 \times 4 + 1) \times (4-1) \times (4 \times 2 - 1) \times (4 \times 3 - 1) \times (4 \times 4 - 1) \times (4 \times 5 - 1) \\ &= (4a + 1)(4b - 1), \text{ where } a, \text{ and } b \text{ are integers.} \\ &= 16ab + 4(b - a) - 1, \text{ the remainder is 3.} \end{aligned}$$

Method 2:  $1 \times 3 \times 5 \times 7 \times 9 \times 11 \times 13 \times 15 \times 17 \times 19 \equiv 1 \cdot (-1) \times 1 \times (-1) \times 1 \times (-1) \times 1 \times (-1) \times 1 \times (-1) \equiv 3 \pmod{4}$

- G9 Given that  $k$ ,  $x_1$  and  $x_2$  are positive integers with  $x_1 < x_2$ . Let  $A$ ,  $B$  and  $C$  be three points on the curve  $y = kx^2$ , with  $x$ -coordinates being  $-x_1$ ,  $x_1$  and  $x_2$  respectively. If the area of  $\triangle ABC$  is 15 square units, find the sum of all possible values of  $k$ .

$\therefore k > 0$ , the parabola opens upwards with following shape:

$$\text{Area of } \triangle ABC = \frac{1}{2} AB \times \text{height}$$

$$\frac{1}{2} (x_1 + x_1) \times (kx_2^2 - kx_1^2) = 15$$

$$kx_1(x_2^2 - x_1^2) = 15$$

$$\text{Possible } k = 1, 3, 5, 15$$


$$\text{When } k = 1, x_1 = 1, x_2 = 4$$

$$\text{When } k = 3, \text{ no solution}$$

$$\text{When } k = 5, x_1 = 1, x_2 = 2$$

$$\text{When } k = 15, \text{ no solution}$$

$$\text{Sum of all possible values of } k = 1 + 5 = 6$$



G10 In Figure 2,  $ABCD$  is rectangular piece of paper with  $AB = 4$  cm and  $BC = 5$  cm. Fold the paper by putting point  $C$  onto  $A$  to create a crease  $EF$ . If  $EF = r$  cm, find the value of  $r$ .

Suppose  $D$  will be folded to a position  $D'$ ,  $\triangle CDE \cong \triangle AD'E$

Let  $O$  be the mid point of  $AC$ .  $AE = CE$ ,  $OE = OE$ ,  $AO = OC$ ,  
 $\triangle AOE \cong \triangle COE$  SSS

$DO = OE$ ,  $\angle ODE = \angle OFB$ ,  $\angle OED = \angle OFB$ ;  $\triangle ODE \cong \triangle OFB$

$CF = BC - BF = AD - DE = AE$

$\therefore AECF$  is a rhombus (4 sides equal)

In  $\triangle ABC$ ,  $AC^2 = 4^2 + 5^2$  (Pythagoras' Theorem)

$AC = \sqrt{41}$ ;  $AO = \frac{\sqrt{41}}{2}$  (diagonal of a rhombus)

Let  $BF = DE = t$

$AE = 5 - t = EC = CF = AF$

In  $\triangle ABF$ ,  $4^2 + t^2 = (5 - t)^2 = 25 - 10t + t^2$  (Pythagoras' Theorem)

$t = \frac{9}{10}$ ,  $AF = 5 - \frac{9}{10} = \frac{41}{10}$

In  $\triangle AOF$ ,  $OF^2 = AF^2 - AO^2$  (Pythagoras' Theorem)

$OF = \sqrt{\left(\frac{41}{10}\right)^2 - \left(\frac{\sqrt{41}}{2}\right)^2} = \frac{4\sqrt{41}}{5}$

Method 2 As before,  $AECF$  is a rhombus.  $AC$  and  $EF$  bisect each other at  $O$ .  $AO = \frac{\sqrt{41}}{2}$ .

$\angle EAO = \angle ACB$  (alt.  $\angle$ s  $AD \parallel BC$ )

$\angle AOE = 90^\circ = \angle ABC$  (property of rhombus)

$\tan \angle EAO = \tan \angle ACB$

$\frac{EO}{\frac{\sqrt{41}}{2}} = \frac{4}{5}$

$EO = \frac{2\sqrt{41}}{5}$

$EF = 2EO = \frac{4\sqrt{41}}{5}$

