

Centripetal Force

The Flying Pig

Purpose


To show the net force for a conical pendulum is mv^2/r .

Equipment and Supplies

Flying Pig (or equivalent) available from Arbor Scientific
vertical and horizontal rod
table clamp
stopwatch
meterstick

Discussion


When an object travels at constant speed along a circular path, we say it has uniform circular motion (if its speed were changing, then its motion would not be uniform). Any object moving in uniform circular motion is accelerated toward the center of its circular path. This acceleration is called *centripetal acceleration*, and equals v^2/r , where v is the speed and r the radius of the circular path. Since the net force on any object equals ma , during uniform circular motion the net force, called *centripetal force* equals mv^2/r and is directed toward the center. This is what happens when an object suspended by a string moves in a circular path—a *conical pendulum*. The string of a conical pendulum sweeps out a right-circular cone. In this experiment you will measure the speed of an object that comprises a conical pendulum and show that the net force is mv^2/r .



Pre-Lab Analysis

Step 1: Draw the force vectors (a free-body diagram) that act on a conical pendulum. Ignoring air resistance, note there are only two forces that act on the pendulum bob, in this case the pig. One is mg , the force due to gravity, and the other is string tension, T .

Step 2: With dashed lines, show the horizontal and vertical components of T . Also label the angle θ , between the tension T and the vertical.




$\theta =$ _____

Step 9: Using the values for r and θ you determined in Steps 7 and 8, compute the theoretical speed of the pig using the formula you derived for the speed in Steps 5 and 6. (Remember, this speed is based on $F_{\text{net}} = mv^2/r$.)

$v =$ _____

Step 10: In Step 9 you calculated the *theoretical* speed of the pig. Now let's actually *measure* the speed of the pig and see how these two values for the speed compare. Since the pig flies in a circle, the speed is the circumference ($2\pi r$) divided by the time t for one complete revolution. To make your measurements of t more precise, measure the time it takes the pig to make 10 revolutions—then divide by 10.



$10t =$ _____ s (for ten revolutions)

So for one revolution, $t =$ _____ s

Step 11: Using your measurement of r , compute the speed of the pig:

$v = d/t = 2\pi r/t =$ _____ m/s

Step 12: Compute the percent difference between the value for the speed you computed in Step 9 and measured in Step 11. (Use the calculated speed as the known.)

% difference = _____

Post-Lab Analysis

1. What techniques for measuring r and θ would you recommend for best results?

2. What do you conclude about the magnitude of the string tension compared with the weight of the pig? For uniform circular motion, the tension will always be (less than) (the same as) (greater than) the weight.

3. What do you conclude about the direction of the net force that keeps the flying pig in uniform circular motion?

4. For uniform circular motion, the centripetal force will always be (less than) (the same as) (greater than) the tension in the string.

5. How does the pig overcome air friction?

6. List sources of error.
