



*The Semantics of ...  
Truth Tables*

[Module Home Page](#)

[Title Page](#)



Page **1** of **23**

[Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

## Lecture 12: The Semantics of Propositional Logic

Aims:

- To discuss what we mean by an interpretation;
- To look at the semantics of compound wffs;
- To discuss the relationship with the semantics of English connectives.



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 2 of 23

Back

Full Screen

Close

Quit

## 12.1. The Semantics of Propositional Logic

Wffs represent statements, and these are either true or false.

### 12.1.1. Atomic Wffs

The truth values of atomic wffs (propositional symbols) are *stipulated*. In other words, we do not spend time debating whether a proposition symbol such as  $p$ ,

$$p \triangleq \text{The moon is made of cheese}$$

denotes a true proposition or a false one. Instead, I stipulate what its truth value is. (In fact, a significant part of what we do will involve considering both what happens when  $p$  denotes a true proposition and when  $p$  denotes a false proposition, so these stipulations aren't required too often anyway.) A stipulation of the truth values of atomic wffs is called an *interpretation* of the atomic wffs. Some books use the phrases *truth assignment* or *value assignment* or *truth valuation* instead of 'interpretation'.

Example (in two different notations):


$p$  has truth value **true**

$q$  has truth value **true**

$r$  has truth value **false**

$\mathcal{I}(p)$  is **true**

$\mathcal{I}(q)$  is **true**



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 3 of 23

Back

Full Screen

Close

Quit

$\mathcal{I}(r)$  is **false**

Both of these ways of writing this interpretation are sentences of the metalanguage. Symbols used in these sentences, other than the proposition symbols themselves, are metasymbols. In particular,  $\mathcal{I}$  is a metasymbol, acting as shorthand for the more verbose version. I've tried to emphasise that it is a metasymbol by using a different font. I've also resisted using = instead of 'is'. We're already using = as an object language symbol in arithmetic, so we don't now want to use it here as a metasymbol.

### 12.1.2. Compound Wffs

The truth value of a compound wff is determined by the truth values of its component wffs. So, we must now look at each connective in turn and see how a wff that is formed using that connective receives its truth value from the truth values of the subwffs of the compound.

### 12.1.3. Negation

Given any wff  $W$ , another wff, called the *negation* of  $W$ , can be formed, and is denoted

$\neg W$

This is read as 'not  $W$ '. Some books use

$\sim W$      $W'$      $\overline{W}$     **not**  $W$

For any wff  $W$ ,



$$\mathcal{I}(\neg W) \triangleq \begin{array}{l} \text{if } \mathcal{I}(W) \text{ is } T \\ \text{then } F \\ \text{else } T \end{array}$$

i.e. to determine the truth value of  $\neg W$ , you must determine the truth value of  $W$ , i.e.  $\mathcal{I}(W)$ . Then,  $\mathcal{I}(\neg W)$  is **true** if  $\mathcal{I}(W)$  is **false** and  $\mathcal{I}(\neg W)$  is **false** if  $\mathcal{I}(W)$  is **true**.

The above is a perfectly good specification of the semantics of  $\neg W$ , but a tabular representation is more often used, as it is clearer:

$W$	$\neg W$
<b>true</b>	<b>false</b>
<b>false</b>	<b>true</b>

You can see that this says the same thing as we said above.

$$\begin{array}{l} p_1 \triangleq \text{Paris is in France} \\ p_2 \triangleq 2 + 2 = 5 \\ p_3 \triangleq \text{Clyde is intelligent} \end{array}$$

Consider the interpretation

$$\mathcal{I}(p_1) \text{ is } \mathbf{true}; \quad \mathcal{I}(p_2) \text{ is } \mathbf{false}; \quad \mathcal{I}(p_3) \text{ is } \mathbf{true}$$

then

$$\mathcal{I}(\neg p_1) \text{ is } \mathbf{false}; \quad \mathcal{I}(\neg p_2) \text{ is } \mathbf{true}; \quad \mathcal{I}(\neg p_3) \text{ is } \mathbf{false}$$

In English, sentences are negated using phrases such as 'It is not the case that', or, perhaps, 'It is false that' and 'It is incorrect that'.

So, using the same propositional symbols from the previous overhead, we have:

The Semantics of ...

Truth Tables

Module Home Page

Title Page

◀ ▶

◀ ▶

Page 4 of 23

Back

Full Screen

Close

Quit



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 5 of 23

Back

Full Screen

Close

Quit

$\neg p_1$  'It is not the case that Paris is in France'  
 $\neg p_2$  'It's not the case that  $2 + 2 = 5$ '

There are other forms of negation in English, where individual words and phrases are negated using 'not' or prefixes such as 'un-' or 'dis-'. Sometimes, these can be straightforwardly represented using propositional logic negation, e.g.

$\neg p_1$  'Paris is not in France'  
 $\neg p_2$   $2 + 2 \neq 5$

Others are less clear. Should 'It's not the case that Clyde is intelligent', 'Clyde isn't intelligent', 'Clyde is unintelligent' and 'Clyde is stupid' all be rendered as  $\neg p_3$ ?

Or consider  $p_4 \triangleq$  All elephants are intelligent.

Should 'It's not the case that all elephants are intelligent', 'Not all elephants are intelligent', 'All elephants are not intelligent', 'All elephants are unintelligent' and 'All elephants are stupid' all be rendered by  $\neg p_4$ ?


$q_1 \triangleq$  Clyde likes Flopsy  
 $q_2 \triangleq$  Clyde is happy

Paraphrase into English

$\neg q_1$

Translate into propositional logic

It's not the case that Clyde is happy.  
Clyde isn't happy.  
Clyde is unhappy.  
Clyde is sad.



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 6 of 23

Back

Full Screen

Close

Quit

### 12.1.4. Conjunction

Any two wffs  $W_1$  and  $W_2$  can be combined to form a compound wff, called the *conjunction* of  $W_1$  and  $W_2$ , and denoted

$$W_1 \wedge W_2$$

$W_1$  and  $W_2$  are called the *conjuncts*. Some books use

$$W_1.W_2 \quad W_1 \& W_2 \quad W_1 \text{ and } W_2 \quad W_1 W_2$$

For any wffs  $W_1$  and  $W_2$ ,

$\mathcal{I}(W_1 \wedge W_2) \triangleq$  if  $\mathcal{I}(W_1)$  is **true** and  $\mathcal{I}(W_2)$  is **true**  
then **true**  
else **false**

Again, the tabular presentation is clearer, so this is what you should learn:

$W_1$	$W_2$	$W_1 \wedge W_2$
true	true	true
true	false	false
false	true	false
false	false	false

Note that we need four rows to capture all the possible combinations of truth values for  $W_1$  and  $W_2$ .

$p_1 \triangleq$  Paris is in France  
 $p_2 \triangleq$  Cork is in Ireland  
 $p_3 \triangleq 2 + 2 = 5$



Consider

$\mathcal{I}(p_1)$  is **true**;       $\mathcal{I}(p_2)$  is **true**;       $\mathcal{I}(p_3)$  is **false**

then, e.g.,

$\mathcal{I}(p_1 \wedge p_2)$  is **true**;       $\mathcal{I}(p_1 \wedge p_3)$  is **false**;       $\mathcal{I}(p_3 \wedge p_3)$  is **false**

The logical behaviour of  $\wedge$  is in some accord with that of the English word ‘and’ (when used to join two sentences):

$p_1 \wedge p_3$     ‘Paris is in France and  $2 + 2 = 5$ ’  
                 ‘Both, Paris is in France and  $2 + 2 = 5$ ’

$q_1 \triangleq$     Clyde likes Flopsy  
 $q_2 \triangleq$     Clyde is happy

Paraphrase into English

$q_1 \wedge \neg q_2$

Translate into propositional logic

It’s not the case that both Clyde likes Flopsy and Clyde is happy.

Certain uses of words such as ‘but’, ‘while’, ‘although’, ‘though’ and ‘whereas’ can also be rendered as logical conjunction.

$p_1 \wedge p_2$     ‘Paris is in France but Cork is in Ireland’  
                 ‘Paris is in France whereas Cork is in  
                 Ireland’  
                 ‘Although Paris is in France, Cork is in  
                 Ireland’

The Semantics of ...

Truth Tables

Module Home Page

Title Page

◀ ▶

◀ ▶

Page 7 of 23

Back

Full Screen

Close

Quit



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 8 of 23

Back

Full Screen

Close

Quit

Some of these words, e.g. 'while' also have completely different meanings that concern the timing of events and these are clearly not rendered using  $\wedge$ .

Also,  $\wedge$  clearly fails to capture the *additional* meaning conveyed by the words 'but', 'while', 'although', 'though' and 'whereas', i.e. that the two sentences express contrasting propositions.

Similarly, not all uses of 'and' have their full meaning captured by  $\wedge$ . For example:

'Batman pulled off his socks and jumped into bed.'

'Batman jumped into bed and pulled off his socks.'

In these, an ordering over the two events being described is also conveyed, but will not be conveyed if we translate these sentences using  $\wedge$ .

The word 'and' can also be used to join phrases and words other than full sentences. Sometimes these can still be translated into propositional logic:

'Batman and Robin are superheroes.'

'Batman is a superhero and Robin is a superhero.'

But other times, such a paraphrase is not necessarily possible, e.g.

'Batman and Robin are fun at parties.'

It might be that they're only fun if they are together, so paraphrasing as 'Batman is fun at parties and Robin is fun at parties' is not necessarily correct.





The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 9 of 23

Back

Full Screen

Close

Quit

### 12.1.5. Disjunction

Any two wffs  $W_1$  and  $W_2$  can be combined to form a compound wff, called the *disjunction* of  $W_1$  and  $W_2$ , and denoted

$$W_1 \vee W_2$$

This is read as ' $W_1$  or  $W_2$ ' or ' $W_1$  or  $W_2$  or both' or ' $W_1$  inclusive-or  $W_2$ '.  $W_1$  and  $W_2$  are called the *disjuncts*. Some books use

$$W_1 + W_2 \quad W_1 \text{ or } W_2$$

For any wffs  $W_1$  and  $W_2$ ,

$\mathcal{I}(W_1 \vee W_2) \triangleq$  if  $\mathcal{I}(W_1)$  is **true** or  $\mathcal{I}(W_2)$  is **true** (or both)  
then **true**  
else **false**

More simply...

$W_1$	$W_2$	$W_1 \vee W_2$
<b>true</b>	<b>true</b>	<b>true</b>
<b>true</b>	<b>false</b>	<b>true</b>
<b>false</b>	<b>true</b>	<b>true</b>
<b>false</b>	<b>false</b>	<b>false</b>

$p_1 \triangleq$  Paris is in France  
 $p_2 \triangleq$  Cork is in Ireland  
 $p_3 \triangleq 2 + 2 = 5$



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 10 of 23

Back

Full Screen

Close

Quit

Consider

$\mathcal{I}(p_1)$  is **true**;       $\mathcal{I}(p_2)$  is **false**;       $\mathcal{I}(p_3)$  is **false**

“Hold on!” you cry. “ $p_2$  is **true** in our world!”

“I know,” I reply. “I just wanted to emphasise the point that *I stipulate* the truth values and you have to go along with them.”

then, e.g.

$\mathcal{I}(p_1 \vee p_2)$  is **true**;       $\mathcal{I}(p_2 \vee p_3)$  is **false**;       $\mathcal{I}(p_1 \vee p_1)$  is **true**

The logical behaviour of  $\vee$  is in some accord with that of the English word ‘or’ (when used to join two sentences):

$p_1 \vee p_3$     ‘Paris is in France or  $2 + 2 = 5$  (or both)’


The English word ‘or’ has at least two distinct uses:

‘ $W_1$ or $W_2$ or both’	<i>inclusive-or</i>
‘ $W_1$ or $W_2$ but not both’	<i>exclusive-or</i>

Inclusive-or is the one we have introduced already.

Exclusive-or is also truth-functional, so we could have a connective for it.  $\oplus$  is the symbol that is often used. We would give this connective a semantics just like that of  $\vee$  except that in the first row of the table, where both  $\mathcal{I}(W_1)$  and  $\mathcal{I}(W_2)$  are **true**,  $\mathcal{I}(W_1 \oplus W_2)$  would be **false**, not **true**. Unless otherwise stated, we are using inclusive-or in this module. Exclusive-or can be captured using  $\vee, \wedge$  and  $\neg$  together (see below).

We see again the advantage of using a formal language in making clear an ambiguity in English. Of course, this means that we need to be extra careful when translating sentences



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 11 of 23

Back

Full Screen

Close

Quit

of English into logic: should we use  $\vee$  or  $\oplus$ ? Generally,  $\vee$  will do. However, sentences that begin with the word ‘either’ often signal use of  $\oplus$ . Also, many people would say that the word ‘unless’ (assuming that it is truth-functional at all) translates as  $\oplus$ . For example, ‘We will go to the cinema unless we go for a pizza’ is probably a sentence being used to convey an exclusive-or: if the speaker of the sentence eventually went to the cinema *and* had a pizza (which is perfectly possible, of course) then I would say that s/he had uttered a false statement. Do you agree? (If not, then you should translate this sentence using  $\vee$ .)

$p_1 \vee p_2$  ‘Paris is in France or Cork  
is in Ireland (or both)’

$(p_1 \vee p_2) \wedge \neg(p_1 \wedge p_2)$  ‘Paris is in France or Cork  
is in Ireland (but not both)’

‘*Either* Paris is in France or  
Cork is in Ireland’

‘Paris is in France unless  
Cork is in Ireland’

‘or’, like ‘and’, can be used to join phrases other than sentences. Only some of these uses of ‘or’ can be translated using the disjunction connective.


### 12.1.6. Conditional

Any two wffs  $W_1$  and  $W_2$  can be combined to form a compound wff denoted

$$W_1 \Rightarrow W_2$$

called a *conditional*. This can be read as ‘if  $W_1$  then  $W_2$ ’.  $W_1$  is called the *antecedent* and  $W_2$  is called the *consequent*. Some books use

$$W_1 \rightarrow W_2 \qquad W_1 \supset W_2$$



The Semantics of ...

Truth Tables

Module Home Page

Title Page

◀ ▶

◀ ▶

Page 12 of 23

Back

Full Screen

Close

Quit

instead of  $W_1 \Rightarrow W_2$ .

Note also that many books call this connective *implication* or *material implication*, instead of the 'conditional'. I try to avoid this because it encourages students to ignore the formal semantics (below) and try to 'get by' using informal paraphrases into English, resulting in them being led astray by this somewhat inappropriate paraphrase.

For any wffs  $W_1$  and  $W_2$ ,

$\mathcal{I}(W_1 \Rightarrow W_2) \triangleq$  if  $\mathcal{I}(W_1)$  is **false** or  $\mathcal{I}(W_2)$  is **true**  
then **true**  
else **false**

More simply...

$W_1$	$W_2$	$W_1 \Rightarrow W_2$
true	true	true
true	false	false
false	true	true
false	false	true

I shall attempt an explanation of why this connective is thought to do something like the job of English 'if...then'. Consider

$p \triangleq$  I win the election  
 $q \triangleq$  Taxes will fall

A politician utters 'If I win the election, then taxes will fall', i.e.  $p \Rightarrow q$ . Under what circumstances would this be a lie?



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 13 of 23

Back

Full Screen

Close

Quit

- Suppose s/he wins and taxes fall:

$\mathcal{I}(p)$  is **true**;       $\mathcal{I}(q)$  is **true**;      so  $\mathcal{I}(p \Rightarrow q)$  is **true**

The semantics says s/he was truthful. Quite right.

- Suppose s/he wins and taxes rise or stay the same:

$\mathcal{I}(p)$  is **true**;       $\mathcal{I}(q)$  is **false**;      so  $\mathcal{I}(p \Rightarrow q)$  is **false**

The semantics says s/he lied in this case. Quite right too.

- Suppose s/he loses and taxes fall:

$\mathcal{I}(p)$  is **false**;       $\mathcal{I}(q)$  is **true**;      so  $\mathcal{I}(p \Rightarrow q)$  is **true**

The semantics says s/he was truthful. See below.

- Suppose s/he loses and taxes rise or stay the same:

$\mathcal{I}(p)$  is **false**;       $\mathcal{I}(q)$  is **false**;      so  $\mathcal{I}(p \Rightarrow q)$  is **true**


The semantics says s/he was truthful. See below.

In these last two cases, it would seem very harsh to say that s/he lied, i.e. that  $\mathcal{I}(p \Rightarrow q)$  is **false**. Of course, it also feels a bit bizarre to say that  $\mathcal{I}(p \Rightarrow q)$  is **true**. But every wff is either **true** or **false**. So we need a convention and the convention is that  $\mathcal{I}(p \Rightarrow q)$  is **true** in both these cases.

$p_1 \triangleq$  Paris is in France

$p_2 \triangleq$  Cork is in Ireland

$p_3 \triangleq 2 + 2 = 5$



The Semantics of ...

Truth Tables

Module Home Page

Title Page

◀ ▶

◀ ▶

Page 14 of 23

Back

Full Screen

Close

Quit

The connection between English and this connective is tenuous at the best of times.

$p_1 \Rightarrow p_2$    ‘If Paris is in France then Cork is in Ireland’  
‘Paris is in France implies Cork is in Ireland’ (avoid!!!)  
‘Paris is in France only if Cork is in Ireland’  
‘Cork is in Ireland if Paris is in France’  
‘Paris being in France is a sufficient  
condition for Cork to be in Ireland’  
‘Cork being in Ireland is a necessary  
condition for Paris to be in France’

Certain uses of ‘when’ and ‘whether’ might also translate using the conditional.

In the above, I’ve deliberately avoided more traditional examples where the antecedent and the consequent are related in some *causal* fashion. For example, if  $p_4 \triangleq$  ‘It is raining’ and  $p_5 \triangleq$  ‘I get wet’ then  $p_4 \Rightarrow p_5$  can be paraphrased as ‘If it is raining, then I get wet’. This sounds very sensible. But now, you are almost certainly thinking that  $\Rightarrow$  says that the antecedent *causes* the consequent. But, we have already seen that ‘because’ is not truth-functional, so treating  $\Rightarrow$ , which is supposed to be a truth-functional connective, as having anything to do with causality must be wrong.


### 12.1.7. Biconditional

Any two wffs  $W_1$  and  $W_2$  can be combined to form a compound wff, denoted

$$W_1 \Leftrightarrow W_2$$

which is called a *biconditional*. This is read as ‘ $W_1$  if and only if  $W_2$ ’. Some books use

$$W_1 \leftrightarrow W_2 \quad W_1 \equiv W_2$$



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 15 of 23

Back

Full Screen

Close

Quit

instead of  $W_1 \Leftrightarrow W_2$ . We'll need the symbol  $\equiv$  as part of our metalanguage later, so we don't want to use it for something else here.

Some books call this connective the 'equivalence' connective but that risks confusion with the metalanguage too (as we shall see). And others call it *bi-implication*. But I'm avoiding the word 'implies' altogether.

For any wffs  $W_1$  and  $W_2$ ,

$\mathcal{I}(W_1 \Leftrightarrow W_2) \triangleq$  if  $\mathcal{I}(W_1)$  is the same as  $\mathcal{I}(W_2)$   
then **true**  
else **false**

More simply...

$W_1$	$W_2$	$W_1 \Leftrightarrow W_2$
true	true	true
true	false	false
false	true	false
false	false	true

$p_1 \triangleq$  Paris is in France

$p_2 \triangleq$  Cork is in Ireland

$p_3 \triangleq 2 + 2 = 5$

Consider

$\mathcal{I}(p_1)$  is **true**;       $\mathcal{I}(p_2)$  is **true**;       $\mathcal{I}(p_3)$  is **false**

then, e.g.,

$\mathcal{I}(p_1 \Leftrightarrow p_2)$  is **true**;       $\mathcal{I}(p_1 \Leftrightarrow p_3)$  is **false**;       $\mathcal{I}(p_3 \Leftrightarrow p_3)$  is **false**



The Semantics of ...

Truth Tables

Module Home Page

Title Page

◀ ▶

◀ ▶

Page 16 of 23

Back

Full Screen

Close

Quit

The following show possible English words and phrases that might translate as the biconditional.

$p_1 \Leftrightarrow p_2$  'Paris is in France if and only if  
Cork is in Ireland'  
'Paris being in France is a necessary  
and sufficient condition for Cork  
to be in Ireland'  
'Paris is in France provided Cork is  
in Ireland'  
'Paris is in France exactly if Cork is  
in Ireland'  
Paris is in France just if Cork is in  
Ireland'


Certain uses of 'when' (e.g. 'just when', 'exactly when') may also translate as the biconditional.

In maths textbooks, the English phrase 'if and only if', which sees fairly regular use in the metalanguage, is often abbreviated to 'iff'.

### 12.1.8. Translations of More Complex Sentences

$p_1 \triangleq$  Paris is in France  
 $p_2 \triangleq$  Cork is in Ireland  
 $p_3 \triangleq 2 + 2 = 5$





The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 17 of 23

Back

Full Screen

Close

Quit

$(p_1 \wedge p_2) \Rightarrow \neg p_3$  'If Paris is in France and Cork  
is in Ireland, then  $2 + 2 \neq 5$

But, beware of ambiguity in English compound statements. English has no precedence and associativity rules to sort this out!

$\neg((p_1 \vee p_2) \Rightarrow p_3)$	
$(p_1 \vee p_2) \Rightarrow \neg p_3$	'It is not the case that
$\neg(p_1 \Rightarrow p_2) \vee p_2$	$2 + 2 = 5$ if Paris is in
$(p_1 \Rightarrow \neg p_3) \vee p_2$	France or Cork is in
$\neg((p_1 \Rightarrow p_3) \vee p_2)$	Ireland'



The Semantics of ...

Truth Tables

Module Home Page

Title Page



Page 18 of 23

Back

Full Screen

Close

Quit

## 12.2. Truth Tables

Let's find the truth-values of a few complex wffs. Given this interpretation:

$\mathcal{I}(p_1)$  is **true**;       $\mathcal{I}(p_2)$  is **false**;       $\mathcal{I}(p_3)$  is **true**

What is  $\mathcal{I}(p_1 \wedge (p_2 \Rightarrow p_3))$ ?

Parentheses determine that the first subwff to be evaluated is  $p_2 \Rightarrow p_3$ . Because  $\mathcal{I}(p_2)$  is **false** and  $\mathcal{I}(p_3)$  is **true**,

$\mathcal{I}(p_2 \Rightarrow p_3)$  is **true**

from row three of the table defining the semantics of  $\Rightarrow$ .

And now we know that  $\mathcal{I}(p_1)$  is **true** and  $\mathcal{I}(p_2 \Rightarrow p_3)$  is **true**, we can determine that

$\mathcal{I}(p_1 \wedge (p_2 \Rightarrow p_3))$  is **true**

from the first row of the table defining the semantics of  $\wedge$ .


So the whole wff is true (*for this interpretation of the atomic wffs*): this wff is true *under* this interpretation.

Given a wff,  $W$ , and an interpretation of the wff's atomic wffs,  $\mathcal{I}$ , we say that  $\mathcal{I}$  *satisfies*  $W$  if and only if  $\mathcal{I}(W)$  is **true** i.e. if the wff is true under that interpretation then that interpretation satisfies that wff.

Hence,  $\mathcal{I}$  from above satisfies  $p_1 \wedge (p_2 \Rightarrow p_3)$ .

Let's do it all again for a different interpretation of the atomic wffs:

$\mathcal{I}(p_1)$  is **true**;       $\mathcal{I}(p_2)$  is **true**;       $\mathcal{I}(p_3)$  is **false**



This time, we'll set things out in a tabular fashion, which results in a tidier presentation. We draw a table with a column for each subwff of the wff:

$p_1$	$p_2$	$p_3$	$(p_2 \Rightarrow p_3)$	$p_1 \wedge (p_2 \Rightarrow p_3)$

Then we can fill in the truth values for the atomic wffs (using our new interpretation  $\mathcal{I}$  from above):

$p_1$	$p_2$	$p_3$	$(p_2 \Rightarrow p_3)$	$p_1 \wedge (p_2 \Rightarrow p_3)$
<b>true</b>	<b>true</b>	<b>false</b>		

Now we fill in the next column with reference to the semantics of  $\Rightarrow$ . This time, the second row of the table that defines the semantics of  $\Rightarrow$  applies:

$p_1$	$p_2$	$p_3$	$(p_2 \Rightarrow p_3)$	$p_1 \wedge (p_2 \Rightarrow p_3)$
<b>true</b>	<b>true</b>	<b>false</b>	<b>false</b>	

And now we can fill in the final column using the semantics of  $\wedge$ . Since  $\mathcal{I}(p_1)$  is **true** and  $\mathcal{I}(p_2 \Rightarrow p_3)$  is **false** (from the third column above),  $\mathcal{I}(p_1 \wedge (p_2 \Rightarrow p_3))$  is **false** (second row of the table defining the semantics of  $\wedge$ ):

$p_1$	$p_2$	$p_3$	$(p_2 \Rightarrow p_3)$	$p_1 \wedge (p_2 \Rightarrow p_3)$
<b>true</b>	<b>true</b>	<b>false</b>	<b>false</b>	<b>false</b>

Obviously, the fourth column here is now not really important. It was just part of our 'working'.



So the whole wff is false under this interpretation, i.e. this interpretation does not satisfy  $p_1 \wedge (p_2 \Rightarrow p_3)$ .

We'll do it yet again, with yet another interpretation:

$$\mathcal{I}(p_1) \text{ is false; } \quad \mathcal{I}(p_2) \text{ is false; } \quad \mathcal{I}(p_3) \text{ is false}$$

And we'll illustrate another way of doing the 'working'.

We draw a table as follows:

$p_1$	$p_2$	$p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$
-------	-------	-------	------------------------------------

We use the interpretation above to fill in the truth values for the atomic wffs:

$p_1$	$p_2$	$p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$
false	false	false	

Then we copy these values to columns headed by occurrences of those wffs in the compound wff:

$p_1$	$p_2$	$p_3$	$p_1$	$p_2 \Rightarrow p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$
false	false	false	false	false	false

Now, since parentheses dictate that the  $\Rightarrow$  is evaluated first, we work out  $\mathcal{I}(p_2 \Rightarrow p_3)$  (which, given that  $\mathcal{I}(p_2)$  is **false** and  $\mathcal{I}(p_3)$  is **false**, is **true**), and we place this into the column headed by the  $\Rightarrow$ :

$p_1$	$p_2$	$p_3$	$p_1$	$p_2 \Rightarrow p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$
false	false	false	false	true	false



And now we evaluate the conjunction  $\mathcal{I}(p_1 \wedge (p_2 \Rightarrow p_3))$  (which, given that we now know that  $\mathcal{I}(p_1)$  is **false** and  $\mathcal{I}(p_2 \Rightarrow p_3)$  is **true**, is **false**), and we write this into the column headed by  $\wedge$ :

$p_1$	$p_2$	$p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$			
false	false	false	false	false	false	false
				$\Delta$		

Obviously, the fourth, sixth, seventh and eighth columns here are not really important: they're just 'working'. The 'result' is in the fifth column.


So, the whole wff is false under this interpretation, i.e. this interpretation does not satisfy  $p_1 \wedge (p_2 \Rightarrow p_3)$ .

Most often we need to compute the truth value of a wff under *every possible* interpretation of its atomic wffs. The tabular presentation on the previous slides can be extended to do a neat job of this: we use one row for each interpretation.

If there are  $n$  different proposition symbols in a compound wff, then there are  $2^n$  different interpretations of those atomic wffs.

The wff  $p_1 \wedge (p_2 \Rightarrow p_3)$  has three different proposition symbols, so there are  $2^3 = 8$  possible interpretations, hence our truth table will have 8 rows.

$p_1$	$p_2$	$p_3$	$p_1 \wedge (p_2 \Rightarrow p_3)$			
true	true	true	true			
true	true	false	false			
true	false	true	true			
true	false	false	true			
false	true	true	false			
false	true	false	false			
false	false	true	false			
false	false	false	false			



- Notes:
- 1. The third, second and eighth rows are the three interpretations we looked at previously.
  - 2. In the lectures I will show you a systematic way to make sure that none of the  $2^n$  rows is forgotten.
  - 3. In the truth table above, I haven't shown any of my 'working'. I'll show this in the lecture.

We will complete these truth tables during the lectures.

$p_1$	$p_2$	$(p_1 \wedge p_2) \Leftrightarrow (p_1 \vee p_2)$
true	true	
true	false	
false	true	
false	false	

$p$	$p \vee \neg p$
true	
false	

$p$	$p \wedge \neg p$
true	
false	

Acknowledgements

Clip Art (of head with bomb) licensed from the Clip Art Gallery on DiscoverySchool.com.



## References

*The Semantics of ...  
Truth Tables*

*Module Home Page*

*Title Page*



*Page 23 of 23*

*Back*

*Full Screen*

*Close*

*Quit*