

ON THE TRANSLATIONALLY-INVARIANT SOLUTIONS OF THE MEMBRANE SHAPE EQUATION

VASSIL M. VASSILEV, PETER A. DJONDJOROV and
IVAÑLO M. MLADENOV

*Institute of Mechanics, Bulgarian Academy of Sciences,
Acad. G. Bonchev Str., Bl. 4, 1113 Sofia, Bulgaria*

*Institute of Biophysics, Bulgarian Academy of Sciences,
Acad. G. Bonchev Str., Bl. 21, 1113 Sofia, Bulgaria*

Abstract. The membrane shape equation [7, 8] describes the equilibrium shapes of a biomembrane, assumed as a bilayer of amphiphilic molecules, in terms of the mean and Gaussian curvatures of its middle-surface. In the present work, a new class of translationally-invariant solutions to this equation in elliptic functions is presented which completes the solutions obtained earlier in [4, 5, 9, 10, 11]. With this, all translationally-invariant solutions to the membrane shape equation are determined. Special attention is paid to those translationally-invariant solutions of the membrane shape equation that determine closed cylindrical (tube-like) surfaces (membrane shapes). Several examples of such surfaces are presented.

1. Introduction

Within the framework of the Helfrich spontaneous curvature model [3], the equilibrium shapes of a biomembrane, assumed as a bilayer of amphiphilic molecules (phospholipids, for instance), are described in terms of the mean H and Gaussian K curvatures of its middle-surface \mathcal{S} by the membrane shape equation [7, 8]

$$2k_c\Delta H + k_c(2H + \mathfrak{h})(2H^2 - \mathfrak{h}H - 2K) - 2\lambda H + p = 0 \quad (1)$$

where k_c , \mathfrak{h} and λ are real constants representing the bending rigidity, spontaneous curvature and tensile stress of the membrane, respectively, while p is the osmotic pressure difference between the outer and inner media assumed to be a real constant too. Here, Δ is the Laplace-Beltrami operator on the surface \mathcal{S} .

In a previous study by the present authors (see [10]), it is established that the six-parameter group of motions in the three dimensional Euclidean space is the largest group of point transformations admitted by the membrane shape equation in Mongé representation. In that work, all types of non-equivalent group-invariant solutions of this equation are identified via an optimal system of one-dimensional subalgebras of the symmetry algebra and the corresponding reduced equations are derived. In [10], special attention is paid to the translationally-invariant solutions of the membrane shape equation assuming that the osmotic pressure difference $p \neq 0$ since the case $p = 0$ is thoroughly studied elsewhere (see [4, 5, 9, 11]). All translationally-invariant solutions of the membrane shape equation that are expressed in elementary functions as well as a class of such solutions that are given in terms of elliptic functions are obtained in [10]. The aim of this study is to determine all other translationally-invariant solutions of the membrane shape equation in Mongé representation.

2. Translationally-Invariant Solutions

In [10], it is shown that the translationally-invariant solutions of the membrane shape equation (1) in Mongé representation correspond to cylindrical surfaces in \mathbb{R}^3 whose directrices are plane curves Γ of curvature $\mathbb{k}(s) = 2H(s)$ that satisfies the equation

$$2 \frac{d^2 \mathbb{k}}{ds^2} + \mathbb{k}^3 - \mu \mathbb{k} - \sigma = 0 \quad (2)$$

where

$$\mu = \mathbb{h}^2 + \frac{2\lambda}{k_c}, \quad \sigma = -\frac{2p}{k_c}$$

s being the arc length of the respective curve Γ . The generatrices of the foregoing cylindrical surfaces are perpendicular to the plane the directrices Γ lie in. Once a solution $\mathbb{k}(s)$ of equation (2) is known in an explicit form, it is possible to recover the embedding $s \mapsto (x(s), y(s))$ of corresponding curve Γ in the plane $\mathbb{R}^2(x, y)$ (up to a rigid motion) by solving the system

$$\frac{dx(s)}{ds} \frac{d^2 y(s)}{ds^2} - \frac{d^2 x(s)}{ds^2} \frac{dy(s)}{ds} = \mathbb{k}(s) \quad (3)$$

$$\left(\frac{dx(s)}{ds} \right)^2 + \left(\frac{dy(s)}{ds} \right)^2 = 1.$$

Thus, the main problem to solve is to find the solutions of equation (2).

This equation is studied by Arreaga *et al* [1] with the aim to determine the equilibria of an elastic loop in the plane subject to the constraints of fixed length and fixed enclosed area. In the three dimensional case considered here, each such loop will

determine a directrix Γ of a cylindrical surface that corresponds to a translationally-invariant solution of the membrane shape equation. In [1], the determination of the curvature \mathbb{k} at equilibrium is reduced to the study of the motion of a particle in a quartic potential. Indeed, equation (2) is the Euler-Lagrange equation associated with the functional

$$F(\mathbb{k}) = \int (T - U) ds, \quad T = \frac{1}{2} \left(\frac{d\mathbb{k}}{ds} \right)^2, \quad U = \frac{1}{8} \mathbb{k}^4 - \frac{1}{4} \mu \mathbb{k}^2 - \frac{1}{2} \sigma \mathbb{k}$$

in which \mathbb{k} , T and U can be thought of as the displacement, kinetic energy and potential energy, respectively, of some fictitious particle. In this setting, s plays the role of the time. Using this analogy, the authors succeeded in obtaining a purely geometric construction for determination of the curvature of the loop passing through a given point of the plane without a reference to explicit expressions for the solutions of equation (2).

However, in our opinion, the knowledge of the solutions of this equation in an explicit form is an important and powerful tool in determination of the surfaces that are translationally-invariant solutions of the membrane shape equation. For this reason, the authors determined some explicit solutions in the previous study [10] and complete this problem here.

Evidently, equation (2) admits the one-parameter group of translations of the independent variable s as a variational symmetry group. Hence, by virtue of Noether's theorem, there is a conservation law of density $E = T + U$, further referred to as the total energy, that is a first integral

$$\frac{dE}{ds} = 0 \tag{4}$$

that holds on its smooth solutions. In characteristic form (see [6]), the above conservation law reads

$$\frac{dE}{ds} = \frac{1}{2} \frac{d\mathbb{k}}{ds} \left(2 \frac{d^2\mathbb{k}}{ds^2} + \mathbb{k}^3 - \mu \mathbb{k} - \sigma \right).$$

Therefore, each solution $\mathbb{k} = \mathbb{k}(s)$ of an equation of form (2), which is not identically a constant, corresponds to a certain real value of the total energy E , and satisfies the equation

$$\left(\frac{d\mathbb{k}}{ds} \right)^2 - P(\mathbb{k}) = 0, \quad P(\mathbb{k}) = 2E - \frac{1}{4} \mathbb{k}^4 + \frac{1}{2} \mu \mathbb{k}^2 + \sigma \mathbb{k}. \tag{5}$$

Evidently, bearing in mind that the coefficients of the polynomial $P(\mathbb{k})$ are real numbers, the foregoing value of the total energy E is such that the polynomial $P(\mathbb{k})$ has at least two different real roots, otherwise the function $P(\mathbb{k}(s))$ could not take non-negative values as required by relation (5). In the light of the above, there are only two possibilities for the roots of the polynomial $P(\mathbb{k})$, namely: (I) the

four roots are real; (II) two of the roots are real and the other two are a complex conjugate pair. Now, denoting by α, β, γ and δ the roots of the polynomial $P(\mathbb{k})$ and specializing to the cases, in which $P(\mathbb{k})$ has simple roots, without loss of generality, in the first case one may consider $\alpha < \beta < \gamma < \delta$ and in the second one $-\alpha, \beta \in \mathbb{R}, \alpha < \beta$ and $\gamma, \delta \in \mathbb{C} \setminus \mathbb{R}, \delta = \bar{\gamma}$. Thus, the polynomial $P(\mathbb{k})$ is nonnegative in the intervals $\alpha \leq \mathbb{k} \leq \beta$ and $\gamma \leq \mathbb{k} \leq \delta$, in case (I), and in the interval $\alpha \leq \mathbb{k} \leq \beta$, in case (II).

All solutions of equation (2) corresponding to multiple roots of the polynomial $P(\mathbb{k})$ are expressed in elementary functions and presented in [10]. That is why, only the cases, in which $P(\mathbb{k})$ has simple roots are considered here.

It should be noted that the roots α, β, γ and δ of the polynomial $P(\mathbb{k})$ can be expressed through its coefficients μ, σ and E as follows

$$\begin{aligned}\alpha &= \sqrt{\frac{\Omega}{2}} + \sqrt{\mu + \sigma\sqrt{\frac{2}{\Omega}} - \frac{\Omega}{2}} \\ \beta &= \sqrt{\frac{\Omega}{2}} - \sqrt{\mu + \sigma\sqrt{\frac{2}{\Omega}} - \frac{\Omega}{2}} \\ \gamma &= -\sqrt{\frac{\Omega}{2}} + \sqrt{\mu - \sigma\sqrt{\frac{2}{\Omega}} - \frac{\Omega}{2}} \\ \delta &= -\sqrt{\frac{\Omega}{2}} - \sqrt{\mu - \sigma\sqrt{\frac{2}{\Omega}} - \frac{\Omega}{2}}\end{aligned}$$

where

$$\begin{aligned}\Omega &= \frac{(\mu + \sqrt[3]{\kappa})^2 - 2^3 3E}{3\sqrt[3]{\kappa}} \\ \kappa &= 3(3^2\sigma^2 + \sqrt{\chi}) - \mu(\mu^2 + 2^3 3^2 E) \\ \chi &= 3(2^3 E((\mu^2 + 8E)^2 - 3^2 2\mu\sigma^2) - \sigma^2(2\mu^3 - 3^3\sigma^2)).\end{aligned}$$

By Vieta's formulas we also have

$$\begin{aligned}\mu &= -\frac{1}{2}(\alpha\beta + \alpha\gamma + \alpha\delta + \beta\gamma + \beta\delta + \gamma\delta) \\ \sigma &= \frac{1}{4}(\alpha\beta\gamma + \alpha\beta\delta + \alpha\gamma\delta + \beta\gamma\delta) \\ E &= -\frac{1}{8}\alpha\beta\gamma\delta.\end{aligned}\tag{6}$$

and

$$\alpha + \beta + \gamma + \delta = 0.\tag{7}$$

due to the absence of a term with \mathbb{k}^3 in the polynomial $P(\mathbb{k})$. The condition $\sigma \neq 0$ implies

$$\alpha + \beta \neq 0, \quad \alpha + \gamma \neq 0, \quad \beta + \gamma \neq 0.\tag{8}$$

3. New Explicit Solutions

Explicit expressions for the solutions of equation (2) are given in Lemma 1 and Lemma 2 for cases (I) and (II), respectively. Lemma 3 shows that any other periodic solution to equation (2) coincides (up to a shift of the independent variable) with one of these solutions.

Lemma 1. *Given μ and σ , let E be such that the roots α , β , γ and δ of the corresponding polynomial $P(\mathbb{k})$ are real numbers ordered as follows: $\alpha < \beta < \gamma < \delta$. Consider the functions*

$$\mathbb{k}(s) = \delta - \frac{(\delta - \alpha)(\delta - \beta)}{(\delta - \beta) + (\beta - \alpha) \operatorname{sn}(us, k)^2} \quad (9)$$

$$\mathbb{k}(s) = \beta + \frac{(\gamma - \beta)(\delta - \beta)}{(\delta - \beta) - (\delta - \gamma) \operatorname{sn}(us, k)^2} \quad (10)$$

of the real variable s in which the parameter u and the elliptic module k are given by the formulae

$$u = \frac{1}{4} \sqrt{(\gamma - \alpha)(\delta - \beta)}, \quad k = \sqrt{\frac{(\beta - \alpha)(\delta - \gamma)}{(\gamma - \alpha)(\delta - \beta)}}. \quad (11)$$

Then, both functions (9) and (10) are real-valued, they are periodic with period $(2/u)K(k)$ and satisfy equation (2).

Proof. It is easy to see that the condition $\alpha < \beta < \gamma < \delta \in \mathbb{R}$ and expressions (11) imply $u \in \mathbb{R}$ and $0 < k < 1$. Therefore, both functions (9) and (10) are real-valued. Evidently, these functions are periodic due to the fact that the function $\operatorname{sn}(us, k)^2$ is periodic with period $(2/u)K(k)$. Finally, substituting each of the above functions into equation (2), one can easily verify that they satisfy it.

Lemma 2. *Given μ and σ , let E be such that two roots, α and β , of the corresponding polynomial $P(\mathbb{k})$ are real numbers ordered as follows: $\alpha < \beta$, and the other two roots, γ and δ , are a complex conjugate pair, that is $\delta = \bar{\gamma}$. Consider the function*

$$\mathbb{k}(s) = \frac{(A\beta + B\alpha) + (A\beta - B\alpha) \operatorname{cn}(us, k)}{(A + B) + (A - B) \operatorname{cn}(us, k)} \quad (12)$$

of the real variable s in which the parameters A , B , u and the elliptic module k are given by the formulae

$$A = \sqrt{4\eta^2 + (3\alpha + \beta)^2}, \quad B = \sqrt{4\eta^2 + (\alpha + 3\beta)^2}. \quad (13)$$

$$u = \frac{1}{4} \sqrt{AB}, \quad k = \sqrt{\frac{1}{2} - \frac{4\eta^2 + (3\alpha + \beta)(\alpha + 3\beta)}{2AB}} \quad (14)$$

where $\eta = (\gamma - \bar{\gamma})/2i$. Then, this function takes real values, it is periodic with period $(4/u) K(k)$ and satisfies equation (2).

Proof. Evidently, in this case, the condition $\alpha < \beta \in \mathbb{R}$ and expressions (14) and (13) imply $u \in \mathbb{R}$ and $0 < k < 1$. Hence, function (12) is real-valued. Obviously, this function is periodic because the function $\text{cn}(us, k)$ is periodic with period $(4/u) K(k)$. Finally, substituting the above function into equation (2), one can easily verify that the latter equation is satisfied.

Lemma 3. Given μ, σ and E , let $\mathbb{k}_i(s)$ ($i = 1, 2$) be two periodic real-valued functions of the real variable s with periods T_i , respectively. Let

$$\frac{d\mathbb{k}_i(s)}{ds} = \sqrt{P(\mathbb{k}_i(s))}, \quad s \in I_i \equiv \left[0, \frac{T_i}{2}\right]$$

and $a = \mathbb{k}_1(0) = \mathbb{k}_2(0)$ is the minimum value of both functions. Then,

$$\mathbb{k}_1(s) = \mathbb{k}_2(s).$$

Proof. Obviously the functions $\mathbb{k}_i(s)$ are invertible for $s \in I_i$, respectively. Let us denote the corresponding inverse functions by $\mathbb{k}_i^{-1}(\mathbb{k})$. Then, differentiating the relations

$$s = \mathbb{k}_i^{-1}(\mathbb{k}) = \mathbb{k}_i^{-1}(\mathbb{k}_i(s))$$

one can see that for each $\mathbb{k} \in (a, b)$ where $b = \min(\mathbb{k}_1(\frac{T_1}{2}), \mathbb{k}_2(\frac{T_2}{2}))$

$$\frac{d\mathbb{k}_i^{-1}(\mathbb{k})}{d\mathbb{k}} = \frac{1}{\sqrt{P(\mathbb{k})}}.$$

Thus, for $\mathbb{k} \in (a, b)$

$$\frac{d}{d\mathbb{k}}(\mathbb{k}_1^{-1}(\mathbb{k}) - \mathbb{k}_2^{-1}(\mathbb{k})) = 0$$

and therefore there exists a real constant s_0 such that

$$\mathbb{k}_2^{-1}(\mathbb{k}) = \mathbb{k}_1^{-1}(\mathbb{k}) + s_0.$$

So,

$$\mathbb{k}_2^{-1}(\mathbb{k}_1(s)) = \mathbb{k}_1^{-1}(\mathbb{k}_1(s)) + s_0 = s + s_0, \quad s \in \min(I_1, I_2)$$

and hence

$$\mathbb{k}_1(s) = \mathbb{k}_2(s + s_0), \quad s \in \min(I_1, I_2).$$

For $s = 0$ this implies $\mathbb{k}_2(s_0) = \mathbb{k}_1(0) = \mathbb{k}_2(0)$ which means that s_0 is a period of the function $\mathbb{k}_2(s)$ and hence the above relation reads

$$\mathbb{k}_1(s) = \mathbb{k}_2(s), \quad s \in \mathbb{R}$$

which completes the proof.

4. Curves and Membrane Shapes

Now, having obtained the solutions of equation (2) in explicit form, one can proceed with constructing the corresponding curves Γ in the plane $\mathbb{R}^2(x, y)$ by solving system (3). Thus, given μ, σ and a solution $\mathbb{k} = \mathbb{k}(s)$ of the corresponding equation (2), without loss of generality, one can represent system (3) in the form

$$\varphi(s) = \int \mathbb{k}(s) ds, \quad \frac{dx(s)}{ds} = \cos(\varphi(s)), \quad \frac{dy(s)}{ds} = \sin(\varphi(s)). \quad (15)$$

Then, using the results presented in [1, 2], which can be cast in the form

$$\begin{aligned} x(s) \frac{dx(s)}{ds} + y(s) \frac{dy(s)}{ds} &= \frac{2}{\sigma} \frac{d\mathbb{k}(s)}{ds} \\ x(s) \frac{dy(s)}{ds} - y(s) \frac{dx(s)}{ds} &= \frac{1}{\sigma} (\mathbb{k}(s)^2 - \mu) \end{aligned}$$

and taking into account relations (15), to write down the explicit expressions

$$\begin{aligned} x(s) &= \frac{2}{\sigma} \frac{d\mathbb{k}(s)}{ds} \cos(\varphi(s)) + \frac{1}{\sigma} (\mathbb{k}(s)^2 - \mu) \sin(\varphi(s)) \\ y(s) &= \frac{2}{\sigma} \frac{d\mathbb{k}(s)}{ds} \sin(\varphi(s)) - \frac{1}{\sigma} (\mathbb{k}(s)^2 - \mu) \cos(\varphi(s)) \end{aligned} \quad (16)$$

for the components of the position vector of the corresponding curve Γ . Computing the integral in (15) one obtains

$$\varphi(s) = \delta s - \frac{\delta - \alpha}{u} \Pi \left(\frac{\beta - \alpha}{\beta - \delta}, \text{am}(us, k), k \right) \quad (17)$$

$$\varphi(s) = \beta s - \frac{\beta - \gamma}{u} \Pi \left(\frac{\delta - \gamma}{\delta - \beta}, \text{am}(us, k), k \right) \quad (18)$$

$$\begin{aligned} \varphi(s) &= \frac{A\beta - B\alpha}{A - B} s + \frac{(A + B)(\alpha - \beta)}{2u(A - B)} \Pi \left(-\frac{(A - B)^2}{4AB}, \text{am}(us, k), k \right) - \\ &\quad - \frac{\alpha - \beta}{2u\sqrt{k + \frac{(A - B)^2}{4AB}}} \arctan \left(\sqrt{k + \frac{(A - B)^2}{4AB}} \frac{\text{sn}(us, k)}{\text{dn}(us, k)} \right) \end{aligned} \quad (19)$$

for the solutions of form (9), (10) and (12), respectively. Thus, given a solution of form (9) or (10), one can draw the corresponding curve substituting the respective angle (17) or (18) in the expressions for the components of the position vector (16). Closed curves are achieved if there exist integers m and n such that

$$\begin{aligned} \frac{\delta}{u} K(k) + \frac{\alpha - \delta}{u} \Pi \left(\frac{\alpha - \beta}{\delta - \beta}, k \right) &= \frac{\pi m}{n} \\ \frac{\beta}{u} K(k) + \frac{\gamma - \beta}{u} \Pi \left(\frac{\gamma - \delta}{\beta - \delta}, k \right) &= \frac{\pi m}{n} \end{aligned}$$




Figure 1. Closed self-intersecting curves and cylindrical surfaces obtained by the solution (9) corresponding to $\mu = 11.82$, $\sigma = -13.3$.

$$\frac{2(A\beta - B\alpha)}{u(A - B)}K(k) + \frac{(A + B)(\alpha - \beta)}{u(A - B)}\Pi\left(-\frac{(A - B)^2}{4AB}, k\right) = \frac{\pi m}{n}$$

for the solutions of form (9), (10) and (12), respectively.

Several examples of closed curves Γ corresponding to solutions of form (9) and (10) of equation (2) with coefficients $\mu = 11.82$ and $\sigma = -13.3$ are presented in Fig. 1 and Fig. 2. It is worthy to underline that in this case, two distinct curves correspond to the same value of the total energy E .

Other examples of closed curves Γ corresponding to solutions of form (12) of equation (2) with coefficients $\mu = -1$ and $\sigma = 1/2$ are presented in Fig. 3.

Acknowledgements

This research is partially supported by the Bulgarian National Science Foundation under a grant K-1531/2005 and the contract # 23/2006 between Bulgarian and Polish Academies of Sciences.

References

- [1] Arreaga G., Capovilla R., Chryssomalakos C. and Guven J., *Area-Constrained Planar Elastica*, Phys. Rev. E **65** (2002) 031801-1-14.




Figure 2. Closed self-intersecting curves and cylindrical surfaces obtained by the solution (10) corresponding to $\mu = 11.82$, $\sigma = -13.3$.




Figure 3. Closed non-self-intersecting curves and cylindrical surfaces obtained by the solution (12) corresponding to $\mu = -1$, $\sigma = 1/2$.

- [2] Guven J., *Laplace pressure as a surface stress in fluid vesicles*, J. Phys. A: Math. Gen. **39** (2006) 3771-3785.
- [3] Helfrich W., *Elastic Properties of Lipid Bilayers: Theory and Possible Experiments*, Z. Naturforsch **28c** (1973) 693-703.
- [4] Ivey T. and Singer D., *Knot Types, Homotopies and Stability of Closed Elastic Rods*, Proc. London Math. Soc. **79** (1999) 429-450.
- [5] Langer J. and Singer D., *Knotted Elastic Curves in \mathbb{R}^3* , J. London Math. Soc. **30** (1984) 512-520.

- [6] Olver P., *Applications of Lie Groups to Differential Equations*, Second Edition, Graduate Texts in Mathematics **107**, Springer, New York, 1993.
- [7] Ou-Yang Z.-C. and Helfrich W., *Instability and Deformation of a Spherical Vesicle by Pressure*, Phys. Rev. Lett. **59** (1987) 2486–2488.
- [8] Ou-Yang Z.-C. and Helfrich W., *Bending Energy of Vesicle Membranes: General Expressions for the First, Second, and Third Variation of the Shape Energy and Applications to Spheres and Cylinders*, Phys. Rev. A **39** (1989) 5280–5288.
- [9] Quan-Hui Liu and Shau-Guang Zhang, *Cylindrical configurations of classical string with rigidity*, J. Phys. A **32** (1999) 5493–5499.
- [10] Vassilev V., Djondjorov P. and Mladenov I., *Symmetry Groups, Conservation Laws and Group-Invariant Solutions the Membrane Shape Equation*, In: Geometry, Integrability and Quantization, I. Mladenov and M. de León (Eds), SOFTEX, Sofia 2006, pp 265–279.
- [11] Zhang Shao-Guang and Ou-Yang Z.-C., *Periodic cylindrical surface solution for fluid bilayer membranes*, Phys. Rev. E **53** (1996) 4206–4208.